

Prescribing of Stoma Discharge Solidifying Agents Position Statement

Hertfordshire and West Essex ICB recommends:

1st Line: Trio Pearls® sachets, maximum monthly quantity of 200 per patient (2 boxes of 100 sachets)*

2nd Line: Respond Absorb® Odour Neutralising Absorbent Gel sachets, maximum monthly quantity of 180 per patient (6 boxes of 30 sachets)*

Quantity required may be considerably more if used for a **fistula. This should be communicated by the local stoma care nurse.*

Recommendations

- For all patients, **Trio Pearls® sachets** should be prescribed as first line option. Maximum monthly quantity of **200** per patient to be prescribed (2 boxes of 100 sachets).
- **Respond Absorb® Odour Neutralising Absorbent sachets** should be prescribed as second line option. Maximum monthly quantity of **180** per patient to be prescribed (6 boxes of 30 strips).
- If ordering in excess of recommended quantities, patients should be referred to their local stoma care nurse for review.
- Authorisation for discharge solidifying agents must come from a stoma care nurse.
- Discharge solidifying agents should not be used for urostomy patients.

Rationale

- Stoma bags may be emptied, on average, up to six times a day.
- Using more than the recommended amount could indicate a stoma related issue that requires a clinical review.
- If used for a urostomy, discharge solidifying agents will block the drainable tap outlet on the device, therefore should not be used for these patients.

Background

- Discharge solidifying agents are added to the bottom of an empty bag to absorb moisture from stoma output, reducing risk of leakage and the need for more frequent bag changes and emptying.
- Discharge solidifying agents should only be used for stoma patients with high watery output.
- Using discharge solidifying agents may mask causes of a high output stoma. Therefore, dietary measures and drug treatment strategies should be addressed before these agents are prescribed. This should be done by the stoma care nurse prior to requesting prescriptions.



Stoma Care Nurse Team Contact Details:

SWH: Dacorum, Watford and St Albans & Harpenden Localities	Telephone: 01923 217489 (Monday-Friday, 8am-4pm)	Email: westherts.stomacare@nhs.net
SWH: Hertsmere Locality	Telephone: 07901 118827 (Monday-Friday, 8:30am-4:30pm)	Email: nclibc.stomanurse@nhs.net
ENH: Stoma office for hospital and community care service, Lister hospital, Stevenage	Telephone: 01438 284133 (Monday-Friday, 7am-4.30pm)	Email: stomacare.enh-tr@nhs.net
West Essex: Stoma care nursing team	Telephone: 01279 827072 (Monday-Friday, 8am-4pm)	Email: paht.stomacare@nhs.net

References:

1. PrescQIPP Stoma and Continence: Adhesive Removers 2.0 (March 2018)
2. Stoma Toolkit Adults, February 2018; NHS Birmingham
3. Stoma Care Products Formulary Guide, September 2011; NHS Buckinghamshire

Version	6
Developed by	Pharmacy and Medicines Optimisation Team, Hertfordshire and West Essex Integrated Care Board Stoma Care Clinical Nurse Specialists, West Hertfordshire Teaching Hospitals NHS Trust, East and North Hertfordshire NHS Trust and The Princess Alexandra Hospital NHS Trust.
Date ratified	V1 July 2020 (Medicines Optimisation Clinical Leads Group). V2 July 2021 (Medicines Optimisation Clinical Leads Group). V3 July 2022 (Medicines Optimisation Clinical Leads group). V4 May 2023 (Medicines Optimisation & Prescribing Delivery and Implementation Group). V5 Updated SWH Hertsmere contact information. V6 May 2024 (Medicines Optimisation & Prescribing Delivery and Implementation Group)
Review date	The recommendation is based upon the evidence available at the time of publication. This recommendation will be reviewed upon request in the light of new evidence becoming available.

