



Primary Care	USCP	Primary or secondary care responsibility	<i>sHRT: sequential</i> <i>ET: endometrial thickness</i>	<i>ccHRT: continuous combined</i> <i>TVS: transvaginal ultrasound</i> <i>USCP: urgent suspicion of cancer pathway</i>
---------------------	-------------	---	---	---

MAJOR risk factors for endometrial cancer

- BMI ≥ 40
- Genetic predisposition (Lynch / Cowden syndrome)
- Estrogen-only HRT for > 6 months in women with a uterus
- Tricycling HRT (quarterly progestogen) for > 12 months
- Prolonged sHRT regimen: use for more than 5 years when started in women aged ≥ 45
- 12 months or more of using norethisterone or medroxyprogesterone acetate for < 10 days / month or, micronised progesterone for < 12 days / month, as part of a sequential regimen

MINOR risk factors for endometrial cancer

- BMI 30-39
- Unopposed estrogen > 3 months but < 6 months
- Tricycling HRT (quarterly progestogen) for > 6 but < 12 months
- > 6 months but < 12 months of using norethisterone or medroxyprogesterone acetate for < 10 days / month or, micronised progesterone for < 12 days / month, as part of a sequential regimen
- Where the progestogen dose is not in proportion to the estrogen dose for > 12 months (including expired 52 mg LNG-IUD)
- Anovulatory cycles, such as in Polycystic ovarian syndrome
- Diabetes